


p-Ethylbenzaldehyde

| | | | |
|-------------------|---|---------------------------|----------------------------------|
| CAS N°: | 4748-78-1 | Empirical formula: | C ₉ H ₁₀ O |
| Structure: |  | | |
| Synonyms: | 4-Ethylbenzaldehyde Benzaldehyde, 4-ethyl (CAS) | | |

| | | | | |
|-----------------|------------------------|------------------------------------|-----------------|--|
| History: | Initial reviews: | New Standard | | |
| | Current revision date: | June 2013 | | |
| | Implementation date: | For new submissions*: | August 10, 2013 | |
| | | For existing fragrance compounds*: | August 10, 2014 | |
| | Next review date | 2018 | | |

* This date applies to the supply of fragrance compounds (formulas) only, not to the finished products in the marketplace.

RECOMMENDATION:

RESTRICTED

RESTRICTIONS:

| Limits in the finished product: | | | |
|--|--------|-------------|------------------|
| Category 1 See Note box (1) | 0.03 % | Category 7 | 0.08 % |
| Category 2 | 0.04 % | Category 8 | 1.11 % |
| Category 3 | 0.17 % | Category 9 | 5.00 % |
| Category 4 | 0.50 % | Category 10 | 2.50 % |
| Category 5 | 0.26 % | Category 11 | See Note box (2) |
| Category 6 | 0.80 % | | |
| Note box: | | | |
| (1) See the IFRA Code of Practice (Appendix 8, Introduction to the IFRA Standards) regarding the Note on Oral Care Products and other products with the potential of ingestion. (2) Category 11 includes all non-skin contact or incidental skin contact products. Due to the negligible skin contact from these types of products there is no justification for a restriction of the concentration of this fragrance ingredient in the finished product. | | | |
| Fragrance material specifications: | | N/A | |

CONTRIBUTION FROM OTHER SOURCES:

None to consider.

p-Ethylbenzaldehyde

CRITICAL EFFECT:
SENSITIZATION
RIFM SUMMARIES:

Cuminaldehyde has been designated as a read across material for p-Ethylbenzaldehyde for the sensitization endpoint. As such, the data for Cuminaldehyde are below:

| LLNA weighted mean EC3 values ($\mu\text{g}/\text{cm}^2$) [no. studies] | Potency Classification Based on Animal Data ¹ | Human Data | | | WoE NESIL ³ ($\mu\text{g}/\text{cm}^2$) |
|---|--|---|--|---|--|
| | | NOEL – HRIPT (induction) ($\mu\text{g}/\text{cm}^2$) | NOEL – HMT (induction) ($\mu\text{g}/\text{cm}^2$) | LOEL ² (induction) ($\mu\text{g}/\text{cm}^2$) | |
| > 2500 [1] ⁴ | Weak | 1181 | 2760 | NA | 1100 ⁵ |

All data in this Table are available from RIFM and are listed in the RIFM Database.

NOEL = No observed effect level; HRIPT = Human Repeat Insult Patch Test; HMT = Human Maximization Test; LOEL = lowest observed effect level; NA = Not Available.

¹Based on animal data using classification defined in ECETOC, Technical Report No. 87, 2003.

²Data derived from HRIPT or HMT.

³WoE NESIL limited to two significant figures.

⁴EC3 value from one LLNA, not the mean.

REXPAN RATIONALE / CONCLUSION:

The RIFM Expert Panel reviewed the critical effect data for Cuminaldehyde and p-Ethylbenzaldehyde and, based on the weight of evidence, established the No Expected Sensitization Induction Level (NESIL) as 1100 $\mu\text{g}/\text{cm}^2$ based on the critical sensitization data on Cuminaldehyde by read-across. They recommend the limits for the 11 different product categories, which are the acceptable use levels of p-Ethylbenzaldehyde in the various product categories. These were derived from the application of the exposure-based quantitative risk assessment approach for fragrance ingredients, which is detailed in the publication by Api *et al.*, 2008.

REFERENCES:

Api AM, Basketter DA, Cadby PA, Cano M-F, Ellis G, Gerberick G, et al. Dermal Sensitization Quantitative Risk Assessment (QRA) For Fragrance Ingredients. *Regulatory Toxicology and Pharmacology* 2008;52(1): 3-23.

RIFM (Research Institute for Fragrance Materials, Inc.), 1972. Maximization test. RIFM report number 1804, November 22. (RIFM, Woodcliff Lake, NJ, USA).

RIFM (Research Institute for Fragrance Materials, Inc.), 1975. Maximization test. RIFM report number 1804, March 27a. (RIFM, Woodcliff Lake, NJ, USA).

RIFM (Research Institute for Fragrance Materials, Inc.), 1973. Maximization test. RIFM report number 1804, November 11. (RIFM, Woodcliff Lake, NJ, USA).

RIFM (Research Institute for Fragrance Materials, Inc.), 2012. Repeat Insult Patch Test. Draft RIFM Report number 63810. (RIFM, Woodcliff Lake, NJ, USA).

RIFM (Research Institute for Fragrance Materials, Inc.) 2012. Local Lymph Node Assay. Draft RIFM Report number 63814. (RIFM, Woodcliff Lake, NJ, USA).