


2-Phenylpropionaldehyde

CAS N°:	93-53-8	Empirical formula: Structure:	$C_9H_{10}O$ 
Synonyms:	Benzeneacetaldehyde, α-methyl-Hydratropaldehyde Hydratropic aldehyde α-Methylphenylacetaldehyde α-Methyltolualdehyde 2-Phenylpropanal α-Phenylpropionaldehyde		

History:	Initial reviews:	New Standard		
	Current revision date:	2009		
	Implementation date:	For new submissions*:	August 7, 2009	
		For existing fragrance compounds*:	August 7, 2011	
Next review date	2014			

* This date applies to the supply of fragrance compounds (formulas) only, not to the finished products in the marketplace.

RECOMMENDATION:

RESTRICTED

RESTRICTIONS:

Limits in the finished product:			
Category 1 See Note box (1)	0.01 %	Category 7	0.03 %
Category 2	0.01 %	Category 8	0.40 %
Category 3	0.06 %	Category 9	1.90 %
Category 4	0.17 %	Category 10	2.50 %
Category 5	0.09 %	Category 11	See Note Box (2)
Category 6	0.28 %		
Note box:			
(1) IFRA would recommend that any material used to impart perfume or flavour in products intended for human ingestion should consist of ingredients that are in compliance with appropriate regulations for foods and food flavourings in the countries of planned distribution and, where these are lacking, with the recommendations laid down in the Code of Practice of IOFI (International Organisation of the Flavor Industry - www.iofiorg.org)			
(2) Category 11 includes all non-skin contact or incidental skin contact products. Due to negligible skin contact the concentration of a fragrance ingredient should not exceed the usual concentration of the fragrance compound in the finished product.			
Fragrance material specifications:	N/A		

CONTRIBUTION FROM OTHER SOURCES:

None to consider (see also the note on contributions from other sources in the **introduction to the IFRA Standards**)

2-Phenylpropionaldehyde

CRITICAL EFFECT:
SENSITIZATION
RIFM SUMMARIES:

LLNA weighted mean EC3 values ($\mu\text{g}/\text{cm}^2$) [no. studies]	Potency Classification Based on Animal Data ²	Human Data			WoE NESIL ³ ($\mu\text{g}/\text{cm}^2$)
		NOEL – HRIPT (induction) ($\mu\text{g}/\text{cm}^2$)	NOEL – HMT (induction) ($\mu\text{g}/\text{cm}^2$)	LOEL ¹ (induction) ($\mu\text{g}/\text{cm}^2$)	
1575 [1] ⁴	Weak	388 ⁵	1380	1938	380 ⁶

All data in this Table are available from RIFM and are listed in the RIFM Database.

NOEL = No observed effect level; HRIPT = Human Repeat Insult Patch Test; HMT = Human Maximization Test; LOEL = lowest observed

effect level; NA = Not Available.

¹ Data derived from HRIPT or HMT.

² Based on animal data using classification defined in ECETOC, Technical Report No. 87, 2003.

³ WoE NESIL limited to two significant figures.

⁴ EC3 value from one LLNA, not the mean.

⁵ HRIPT with 38 subjects only; study with 100 subjects not done because the material is used at a low volume (1-5 metric tons)

⁶ WoE NESIL based on limited subject HRIPT which was lower than the default LLNA (1000 $\mu\text{g}/\text{cm}^2$)

REXPAN RATIONALE / CONCLUSION:

The RIFM Expert Panel reviewed the critical effect data for 2-Phenylpropionaldehyde and, based on the weight of evidence, established the No Expected Sensitization Induction Level (NESIL) as 380 mg/cm^2 . They recommend the limits for the 11 different product categories, which are the acceptable use levels of 2-Phenylpropionaldehyde in the various product categories. These were derived from the application of the exposure-based quantitative risk assessment approach for fragrance ingredients, which is detailed in the QRA Expert Group Technical Dossier of June 22, 2006.

REFERENCES:

Patlewicz, G., Roberts, D.W., Walker, J.D., 2003. QSARs for the skin sensitization potential of aldehydes and related compounds. QSAR & Combinatorial Science, 22, 196-203.

RIFM (Research Institute for Fragrance Materials, Inc.), 1964. Human repeated insult patch test. Unpublished study from IFF Inc., 3 April. Report number 51926. (RIFM, Woodcliff Lake, NJ, USA).

RIFM (Research Institute for Fragrance Materials, Inc.), 1971a. Human repeated insult patch test. Unpublished study from IFF Inc., 7 July. Report number 51925. (RIFM, Woodcliff Lake, NJ, USA).

RIFM (Research Institute for Fragrance Materials, Inc.), 1971b. Maximization study with 2-phenylpropionaldehyde. RIFM report number 1805, April 2a. (RIFM, Woodcliff Lake, NJ, USA).