


trans-2-Hexenal dimethyl acetal

CAS-No.:	18318-83-7 The scope of this Standard includes, but is not limited to the CAS number(s) indicated above; any other CAS number(s) used to identify this fragrance ingredient should be considered in scope as well.	Molecular formula:	C ₈ H ₁₆ O ₂
		Structure:	
Synonyms:	1,1-Dimethoxy-trans-2-hexene 2-Hexene, 1,1-dimethoxy-, (2E)-		

History:	Publication date:	2006 (Amendment 40)	Previous Publications:	1985 1989 2002
-----------------	-------------------	---------------------	------------------------	----------------------

Implementation dates:	For new submissions*:	Not applicable.
	For existing fragrance compounds*:	Not applicable.
*These dates apply to the supply of fragrance mixtures (formulas) only, not to the finished consumer products in the marketplace.		

RECOMMENDATION:
PROHIBITION
FRAGRANCE INGREDIENT PROHIBITION:

trans-2-Hexenal dimethyl acetal should not be used as a fragrance ingredient.

CONTRIBUTIONS FROM OTHER SOURCES:
NONE TO CONSIDER (SEE ALSO THE SECTION ON CONTRIBUTIONS FROM OTHER SOURCES IN CHAPTER 1 OF THE GUIDANCE FOR THE USE OF IFRA STANDARDS)

trans-2-Hexenal dimethyl acetal

INTRINSIC PROPERTY MANAGEMENT:	DRIVING RISK	DERMAL SENSITIZATION
---------------------------------------	---------------------	-----------------------------

EXPERT PANEL FOR FRAGRANCE SAFETY RATIONALE / CONCLUSION:

The Expert Panel for Fragrance Safety reviewed all the available data for trans-2-Hexenal dimethyl acetal and recommends not to use trans-2-Hexenal dimethyl acetal as or in fragrance ingredients in any finished product application.

REFERENCES:

The IFRA Standard on trans-2-Hexenal dimethyl acetal is based on at least one of the following publications:

- The RIFM Safety Assessment on trans-2-Hexenal dimethyl acetal is available at the RIFM Safety Assessment Sheet Database: <http://fragrancematerialsafetyresource.elsevier.com/>.
- Api A.M., Belsito D., Bruze M., Cadby P., Calow P., Dagli M. L., Dekant W., Dent M., Ellis G., Fryer A. D., Fukayama M., Griem P., Hickey C., Kromidas L., Lalko J., Liebler D.C., Miyachi Y., Politano V.T., Renskers K., Ritacco G., Salvito D., Schultz T.W., Sipes I. G., Smith B., Vitale D., Wilcox D.K. (2015). Criteria for the Research Institute for Fragrance Materials, Inc. (RIFM) safety evaluation process for fragrance ingredients. *Food Chem Toxicol.* 2015 Aug;82 Suppl:S1-S19 (doi: 10.1016/j.fct.2014.11.014). (http://fragrancematerialsafetyresource.elsevier.com/sites/default/files/Criteria_Document_Final.pdf).
- IDEA project (International Dialogue for the Evaluation of Allergens) Final Report on the QRA2: Skin Sensitisation Quantitative Risk Assessment for Fragrance Ingredients, September 30, 2016 (<http://www.ideaproject.info/uploads/Modules/Documents/qra2-dossier-final--september-2016.pdf>).
- Salvito D.T., Senna R. J., Federle T.W. (2002). A framework for prioritizing fragrance materials for aquatic risk assessment. *Environ Toxicol Chem.* 2002;21:1301-1308. (<https://www.ncbi.nlm.nih.gov/pubmed/12069318>).
- Ford, R.A. (1988), *Food and Chemical Toxicology* 26, 347.

Additional information on the application of IFRA Standards is available in the Guidance for the use of IFRA Standards, publicly available at www.ifrafragrance.org.